# **Discharge Instructions for Cataract Surgery**

A surgeon has removed the cloudy lens in your eye and replaced it with a clear manufactured lens. Be sure to have an adult family member or friend drive you home after surgery. Here's what you can expect after surgery, and some tips for a healthy recovery.

#### What to expect

It's normal to have the following:

- Bruised or bloodshot eye for 7 days
- Mild floaters
- Itching and mild pain for several days
- Sensitivity to light
- Scratchy, sandlike feeling in the eye for several days
- Feeling tired, especially during the first 24 hours

### **Activity level**

- Don't drive for 2 days or as instructed by your eye care provider.
- Don't drink alcohol for at least 24 hours.
- Don't bend at the waist to pick up objects or lift anything heavy for 1 to 2 weeks.
- Relax for the first 24 hours after surgery. Watching TV and reading are OK and won't harm your eye.

#### Eye protection

- Don't rub or press on your eye.
- If instructed, wear a bandage over your eye for 2 days and 2 nights.
- If instructed, wear a shield to sleep to protect your eye for 1 week.
- While you're healing, your eye health and vision will be watched. Ask your provider how often you will need to be seen.
- After your eyes have fully healed, your eye care provider can evaluate if you need eyeglasses or contact lenses.

#### Using eye drops

You may be given special eye drops or ointment. Here is one way to use eye drops:

© Krames Page 1 of 2

- Wash your hands before and after using the drops.
- Tilt your head back.
- Pull your bottom eyelid down.
- Squeeze 1 drop into your eye. Don't touch your eye with the bottle tip.
- Close your eyes for a few seconds.
- If you need more than 1 drop, wait at least 5 minutes before adding the next one.

## When to call the healthcare provider

Call your healthcare provider if you have any of these:

- Bleeding or discharge from the eye
- Your vision suddenly gets worse
- Flashes of light or a lot of new floaters
- Pain that doesn't improve with the pain medicine that is advised for you
- Nausea or vomiting
- Chills or fever of 100.4°F (38°C) or higher, or as directed by your provider

© Krames Page 2 of 2